"Products to grow with"

Ca 6% B 2%

CALCIUM/ BORON MIX

FOR COMMERCIAL USE ONLY

PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc . or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

GUARANTEED ANALYSIS:

Calcium (Ca)......6.00% Boron (B).....2.00% Chlorine (CI), not more than..... 11.00% Derived from: Calcium Chloride and Sodium Borate.

WARNING

Overapplication or excessive use of boron, or use on crops other than those listed, can result in boron toxicity. This may cause foliar damage or serious crop injury.

DIRECTIONS FOR USE:

The rate and timing of application will depend on local conditions and should always be made as a result of soil and/or plant tissue analysis.

Broadcast: Severe Deficiency 4-6 qts. per acre, Moderate Deficiency 2-4 gts. per acre, or Maintenance 1-2 gts. per acre.

Banded over the Row: Apply at 2-4 qts. per acre. Repeat as necessary if deficiency persists.

Foliar Application: Apply at 2-4 qts. per acre. Repeat as necessary if deficiency persists. Use sufficient water for thorough coverage. Ground: 1-20 gals. per acre. Air: 3-5 gals. per acre. If less water is used, slight burn may occur.

*Spray either early or later in the day for maximum leaf absorption.

***Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a iar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is

CAUTION: KEEP OUT OF REACH OF CHILDREN

Net Content: Size / Weight

2.5 gl/29.1 lbs 5 gl/58.05lbs 30 gl/348.3 lbs 55 gl/638.55lbs 275 gl/3,192.75 lbs

Manufactured by:



534 CR 529A Lake Panasoffkee, FL 33538 (352)-793-1682 www.uas-cropmaster.com F001142

UNITED AGRICULTURAL SERVICE many other environmentally safe products. NPK/Soil products t expose to sunlight for long periods (77 F). s, seaweed, fish emulsion, nitrogen-fixing baterica, The hot-mixed, homogenized CROPMASTER PLANT s of high-quality oners. These food-grade purity

Storage: Keep container sealed tightly when not of time. Store in temperatures between 5 C (41

nt in use. Do not 1 F) and 25 C (