



GUARANTEED ANALYSIS: Non-Plant Food Ingredients: Fulvic Acid......25.0% F001142

Derived from: North Dakota Leonardite Shale DIRECTIONS FOR USE:

Preparation for Germination (in indoor environment)

First Prepare mix for every 1 gallon of water add 4 oz of Seed and Plant Starter.

- 1. Fill your seedling flat with potting medium mix of your choice
- 2. Moisten with Super Seed Starter mixed with water.
- 3. Plant your seeds approximately 1" apart at a depth of 1/4 to 1/2 inch
- 4. Cover with potting medium
- 5. Place in warm, lighted environment
- 6. Keep moist with daily light watering

After the seedlings begin to germinate, begin fertilizing them with a Seed and Plant Starter mixed with water every other day until they reach desire height. then plant in field.

Outdoor Field Germination

Mix 1.25 gallon of Seed and Plant Starter for each acre, mixed with maximum 30 to 50 gallons of water. Solution is used to moisten area where seed is dropped until seed is totally soaked. Also moisten area at least 3 inches around seed until damp with mix.

***Highly concentrated DO NOT over apply. *** To ensure compatibility with foreign spray mixes, a jar compatibility test is recommended before mixing with other products other than water in the tank. Always fill tank with water first, then begin applying CROPMASTER products while tank is circulating. "NEVER MIX CONCENTRATES DIRECTLY TOGETHER "

CAUTION: KEEP OUT OF REACH OF CHILDREI

Net Content: Size / Weight

5 gl/51.10 lbs 2.5 gl/25.55 lbs

FOR COMMERCIAL USE ONLY PRECAUTIONS

Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eves, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

The liquid analysis results were obtained from a third party laboratory using the Separation and Analysis of Humic Acid and Fulvic Acid components of humates. (Verploegh and Brandvold 1990).

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or it's be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

UAS

Manufactured by:

534 CR 529A Lake Panasoffkee, FL 33538 (352)-793-1682 www.uas-cropmaster.com

ġ

≤

germinate

fa

lster

stronger

seed starting with consistent and tremendous results

Tomato seed germinate

Near 100% of the seed Will germinate Either start seed l indoors ≤. or outdoors in field the 2 leaf stage 2 inches inches within 5 to in 24 and 48 hours germinate

- CORN to 7 days



Dispose of Waste Material according to local, state, and federal environmental laws and regulations.