

GUARANTEED ANALYSIS

 Total Nitrogen (N)
 3.00%

 1.20% Ammoniacal Nitrogen
 .35% Nitrate Nitrogen

 1.45% Urea Nitrogen
 .35%

 Available Phosphate (P2O5)
 18.00%

 Soluble Potash (K2O)
 18.00%

 Derived from: Urea, Potassium Nitrate, Ammonium Hydroxide, and Orthophosphoric Acid.

DIRECTIONS FOR USE

Lawns/Turf: Apply at a rate of 1 quart per 4000 SQ. FT. Set hose end dial at 2oz per gallon. Apply every 3-5 weeks.

Ornamentals: 0.5 to 1oz Super 3-18-18 per gallon of water mix. Mist tops and bottoms of matured plant leaves. Use after blooming stage.

Row Crops : Mix 1-2 quarts Super 3-18-18 with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off. (USE MINIMUM OF 5 GALLONS OF WATER) Apply after Fruit has changed color or is beginning to mature.

Tree Crops : Mix 1 gallon Super 3-18-18 with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off. (USE MINIMUM OF 50 GALLONS OF WATER) Apply after Fruit has changed color or is beginning to mature.

*Can be applied every two weeks until 10 days prior to harvest.

These are general product recommendations, please consult with your sales representative for other applications.

Net Content: Size / Weight

2.5 al/29.3 lbs

Super 3-18-18

NUTRITIONAL SUPPLEMENT FOR FLOWERS, FRUIT AND VEGETABLES

PRODUCT INFORMATION

- * Increase Fruit Set
- * Increase Fruit Size
- * Enhance Bloom
- * Low Salt Index
- * Near Neutral pH
- * Low Impurities

PRECAUTIONS

Avoid getting in eyes or on skin. The use of safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

DISCLAIMER

It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended.

Always read label instructions.

CONDITIONS OF SALE

UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the buyer/end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

FOR COMMERCIAL USE ONLY



534 CR 529A Lake Panasoffkee, FL 33538 (352)-793-1682

www.uas-cropmaster.com F001142 control testing of these aw materials before s manufacturing uality LIQUID mogenized NPK formu control BIOLOGICAL tests to ensure that at all times, process formulation products ulation. N-P-K ಕ go through FORMULATION with as Ve also With these safeguards. a series use only inspectior the Ve of control ₽ end result ie best have മ superior edge over the competition. taken natural r steps to always ឆ the highest quality our nontoxic food grade accomplish the end industry guarantee into the Ŧ product. end e end raw mat resul , م We have added aterials in this pr uniform high quality product also do quality With each step we carefully our quality product. bio-stimulants to roduct. UAS of A checks on this hot mix and 8 idard

Storage: Keep container sealed tightly when not in use. temperatures between 5 C (41 F) and 25 C (77 F).

Do not expose to sunlight for long periods of time.

Store in

Weight CAUTION: KEEP OUT OF REACH OF CHILDREN