



# SA 3-0-6

## DIRECTIONS FOR USE:

### SHAKE BEFORE USING.

Turf: With sufficient water to cover, apply up to 3 oz per 1000 ft2. Start applications in initial growth stage and continue every 3 to 4 weeks. Use monthly to help retain turf health and vigor.

Row Crops: Mix 4-32 ozs. SA 3-0-6 with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off. (Use Minimum of 5 gallons of water). To be used when plants are in growth stage after 2nd leaf set has occurred. Can be applied every 4-6 weeks.

Tree Crops: Apply at a rate of 8-32 ozs. SA 3-0-6 with sufficient water to cover, per acre. Mist tops and bottoms of leaves. Moisten only to the point of run-off. (Use Minimum of 50 gallons of water). Apply during growth stage. Can be applied every 4-6 weeks.

**Dispose of Waste Material** according to local, state, and federal environmental laws and regulations.

**Storage:** Keep container sealed tightly when not in use. Do not expose to sunlight for long periods of time. Store in temperatures between 5°C (41°F) and 25°C (77°F).

**Precautions:** Avoid getting in eyes or on skin. The use of side shield safety glasses and gloves is recommended. Harmful if swallowed. Avoid contact with eyes, skin, and clothing. If on skin, remove contaminated clothing. Wash with large amounts of soap and water. If in eyes, rinse repeatedly with clean water for 15 minutes. If irritation persists, obtain medical attention. See MSDS for further information.

**Lot #:**

**Net Contents: Size / Weight**

2.5 g/23.20 lbs  5 g/46.40 lbs  30 g/278.40 lbs  55 g/510.40 lbs  275 g/2552 lbs

## GUARANTEED ANALYSIS:

Total Nitrogen (N) .....3.00%  
3.00% Urea Nitrogen  
Available Phosphate (P205) .....0.00%  
Soluble Potash (K2O) .....6.00%  
Derived from: Urea and Potassium Citrate

Non-Plant Food Ingredients:

3.50% Salicylic Acid  
Derived from: Salicylic Acid

**Manufactured by:**



**UAS of America, Inc. - 534 CR 529A Lake Panasoffkee,  
FL 33538 - (352)-793-1682 - www.cropmaster.com  
F001142**

## CAUTION: KEEP OUT OF REACH OF CHILDREN

**Conditions of Sale:** UAS of America Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, product use ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of UAS of America Inc. In no case shall UAS of America Inc. or its be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product.

All such risk shall be assumed by the buyer /end user. UAS of America Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

**Disclaimer:** Highly Concentrated, Do Not Over Apply. Never Mix Concentrates Directly together. It is recommended not to use this product with fungicides or pesticides. However, if applied with a fungicide or pesticide, it is the users responsibility to conduct a trial application on the crop(s) to be sprayed with this mix of chemicals to determine compatibility of products and compatibility with crop(s) sprayed. When mixing with other spray materials a jar compatibility test is recommended. Always read label instructions. When adding into the tank: fill tank first with water then add in CropMaster product while tank is circulating.